AMENDMENT TO THE CLAIMS:

Please amend the claims as follows:

1. (Original) A polymer comprising an optionally substituted repeat unit of formula (I):

$$R$$
 R
 R
 R
 R

(I)

wherein each R is the same or different and represents H or an electron withdrawing group; and each R^1 is the same or different and represents a substituent.

- 2. (Currently Amended) A polymer according to claim 1 wherein at least one R¹ is a solubilising solubilizing group.
- 3. (Currently Amended) A polymer according to claim 1 or 2 wherein each R^1 is the same or different and is independently selected from the group consisting of optionally substituted C_{1-20} alkyl, C_{1-20} alkoxy, aryl and heteroaryl groups.
- 4. (Currently Amended) A polymer according to any preceding claim 1 comprising an optionally substituted aryl or heteroaryl second repeat unit.
 - 5. (Currently Amended) A monomer comprising a repeat unit of formula (II):

$$X$$
 R^{1}
 Si
 R^{1}
 (II)

wherein R and R1 are as defined in any one of claims 1-3 each R is the same or different and represents H or an electron withdrawing group; and each R¹ is the same or different and represents a substituent and each X independently represents a polymerisable polymerizable group.

- 6. (Currently Amended) A monomer according to claim 5 wherein each X is the same or different and is selected from the group consisting of boronic acid groups, boronic ester groups, borane groups, and halide functional groups.
- 7. (Currently Amended) A method of forming a polymer comprising the step of polymerizing a monomer according to claim 5 or 6.
- 8. (Currently Amended) A method according to claim 7 wherein each X is the same or different and is a halide functional group, and comprising performing the polymerization polymerization is performed in the presence of a nickel complex catalyst.
- 9. (Currently Amended) A method according to claim 7 comprising the step of polymerizing:
- (a) a monomer of formula (II) wherein each X is a boron the same or different and is a boron derivative functional group selected from [[a]] the group consisting of boronic acid, [[a]] boronic ester esters, and [[a]] borane boranes, and an aromatic monomer having at least two reactive halide functional groups; or

- (b) a monomer of formula (II) wherein each X is the same or different and is a reactive halide functional group, and an aromatic monomer having at least two boron derivative functional group selected from [[a]] the group consisting of boronic acid, [[a]] boronic ester esters, and [[a]] borane boranes; or
- (c) a monomer of formula (II) wherein one X is a reactive halide functional group and the other X is a boron derivative functional group selected from [[a]] the group consisting of boronic acid, [[a]] boronic ester esters, and [[a]] borane boranes,

wherein the reaction mixture comprises a catalytic amount of a palladium catalyst suitable for eatalysing catalyzing the polymerisation polymerization of the aromatic monomers, and a base in an amount sufficient to convert the boron derivative functional groups into boronate anionic groups.

- 10. (Currently Amended) An optical device comprising a polymer according to any one of claims 1-4 claim 1.
- 11. (Currently Amended) An optical device according to claim 10 comprising an anode, a cathode and a layer of the polymer according any one of claims 1-4 located disposed between the anode and the cathode.
- 12. (Currently Amended) An optical device according to claim 11 that is comprising an electroluminescent device.
- 13. (Currently Amended) A switching device comprising a polymer according to any one of claims 1-4 claim 1.

- 14. (Currently Amended) A switching device according to claim 13 that is comprising a thin film transistor.
 - 15. (Currently Amended) An optionally substituted compound of formula (IV):

$$X^1$$
 X^2
 X^2
 X^2
 X^3
 X^4
 X^2
 X^4
 X^2
 X^4

(IV)

wherein R is as defined in any one of claims 1-3 the same or different and represents

H or an electron withdrawing group; each X^1 and each X^2 are the same or different and
represent a leaving group capable of participating in a transmetallation reaction and X^2 has an electronegativity less than that of X^1 .

- 16. (Currently Amended) Preferably, The compound of claim 15, wherein each X^1 and X^2 is independently a halogen.
- 17. (Currently Amended) A method of forming a monomer of formula (VI) from a compound of formula (V) according to the following scheme::


wherein the method comprises reacting the compound of formula (V) with a transmetallating agent followed by reaction with a compound of formula LG-Y-LG, wherein X^1 is a leaving group capable of participating in a transmetallation reaction and R are as defined in claim 15 is H or an electron withdrawing group; each X^3 is the same or different

and represents a leaving group capable of participating in a transmetallation having an electronegativity less than or the same as that of X^1 ; Y represents a divalent residue comprising a backbone of 1-3 atoms; and each LG is the same or different and represents a leaving group.

- 18. (Currently Amended) A method according to claim 17 wherein Y comprises a single atom in its backbone selected from the group consisting of $-CR_{2}^{3}$, $-SiR_{2}^{3}$, $-NR_{3}^{3}$, $-PR_{2}^{3}$, $-GeR_{2}^{3}$, $-SnR_{2}^{3}$, O, and S, wherein R³ is selected from the group consisting of optionally substituted alkyl, alkoxy, aryl, and heteroaryl.
- 19. (Currently Amended) A method according to claim 17 or 18 wherein each X^3 is the same or different and has an electronegativity less than that of X^1 .
- 20. (Currently Amended) A method according to any one of claims 17-19 claim

 17 wherein each LG is the same or different and is a halogen.
- 21. (Currently Amended) A method according to any one of claims 17-20 claim

 17 wherein the transmetallating agent is a compound of formula R⁴-M wherein R⁴ is alkyl or aryl and M is a metal.
- 22. (Original) A polymer comprising an optionally substituted first repeat unit of formula (VII):



(VII)

wherein each R² is the same or different and represents a substituent; the R² groups may be linked to form a ring; and bond (a) is not linked to the 2-position of the repeat unit of formula (VII).

- 23. (Original) A polymer according to claim 22 wherein bond (b) is not bound to the 7-position of the repeat unit of formula (VII).
- 24. (Currently Amended) A polymer according to claim 22 or 23 wherein bond(a) is bound to the 3-position of the repeat unit of formula (VII).
- 25. (Currently Amended) A polymer according to any one of claims 22-24 claim 22 wherein bond (b) is bound to the 6-position of the repeat unit of formula (VII).
- 26. (Currently Amended) A polymer according to any one of claims 22-25 claim

 22 wherein at least one R² is a solubilising group.
- 27. (Currently Amended) A polymer according to any one of claims 22-26 claim 22 wherein each R² is the same or different and is selected from the group consisting of optionally substituted C₁₋₂₀ alkyl, C₁₋₂₀ alkoxy, aryl and heteroaryl, preferably a C4-10 alkyl, more preferably n-hexyl or n-octyl.
- 28. (Currently Amended) A polymer according to any one of claims 22-27 claim

 22 wherein the polymer comprises an optionally substituted aryl or heteroaryl second repeat
 unit.
 - 29. Currently Amended) An optionally substituted monomer of formula (VIII):

(VIII)

wherein each R² is as defined in claim 22, 26, or 27 the same or different and represents a substituent; each X is as defined in claim 5 or 6 independently represents a polymerizable group and at least one X is not linked to the 2-position of the repeat unit of formula (VIII).

30. (Currently Amended) An electroluminescent device comprising an anode, a cathode and an electroluminescent layer located between the anode and cathode wherein the electroluminescent layer comprises a polymeric host material comprising an optionally substituted first repeat unit of formula (IX) and a luminescent dopant

$$R^{1}$$
Si R^{1}

(IX)

wherein R¹ is as defined in any one of claims 1-3 the same or different and represents a substituent.

- 31. (Original) An electroluminescent device according to claim 30 wherein the repeat unit of formula (IX) is linked through its 3- and 6- positions.
- 32. (Original) An electroluminescent device according to claim 30 or 31 wherein the polymeric host material comprises a second repeat unit

- 33. (Original) An electroluminescent device according to any one of claims 30-32 claim 30 wherein the second repeat unit comprises a hole transporting material.
- 34. (Original) An electroluminescent device according to any one of claims 30-33 claim 30 wherein the luminescent dopant is phosphorescent.
- 35. (Original) A method of forming an optionally substituted compound of formula (X) according to the following process:

wherein each R^8 is independently selected from the group consisting of C_{1-20} alkyl and aryl; each R^9 is different from R^8 and is independently selected from the group consisting of C_{1-20} alkyl, aryl and heteroaryl; M^1 is a metal; and Z is a reactive group capable of undergoing reaction with M^1 - R^9 .

- 36. (Original) A method according to claim 35 wherein M¹ is lithium.
- 37. (Currently Amended) A method according to claim 35 wherein R⁸ is methyl.
- 38. (Currently Amended) A method according to claim 35 wherein Z is trialkylsilyl, more preferably trimethylsilyl.

- 39. (Original) A method according to claim 35 wherein, in the case of reaction with M^1 - R^9 , the two groups R^{10} are not linked to form a ring.
 - 40. (New) A polymer according to claim 22, wherein R² is a C₄₋₁₀ alkyl group.
- 41. (New) A polymer according to claim 40, wherein R² is a m-hexyl group or an n-octyl group.
 - 42. (New) As method according to claim 35, wherein Z is trimethylsilyl.